



Performance bound for Approximate Optimistic Policy Iteration

Bruno Scherrer, Christophe Thiery

► To cite this version:

Bruno Scherrer, Christophe Thiery. Performance bound for Approximate Optimistic Policy Iteration. [Technical Report] 2010. inria-00480952

HAL Id: inria-00480952

<https://inria.hal.science/inria-00480952>

Submitted on 5 May 2010

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Performance bound for Approximate Optimistic Policy Iteration

Christophe Thiery, Bruno Scherrer

We provide here a proof of the performance bound theorem published in Thiery and Scherrer (2010). This theorem applies to Least-Squares λ Policy Iteration and more generally approximate, optimistic Policy Iteration algorithms.

Theorem 1 (Performance bound for Approximate Optimistic Policy Iteration)

Let $(\lambda_n)_{n \geq 1}$ be a sequence of positive weights such that $\sum_{n \geq 1} \lambda_n = 1$. Let Q_0 be an arbitrary initialization. We consider an iterative algorithm that generates the sequence $(\pi_k, Q_k)_{k \geq 1}$ with

$$\begin{aligned}\pi_{k+1} &\leftarrow \text{greedy}(Q_k), \\ Q_{k+1} &\leftarrow \sum_{n \geq 1} \lambda_n (B_{\pi_{k+1}})^n Q_k + \epsilon_{k+1}.\end{aligned}$$

ϵ_{k+1} is the approximation error made when estimating the next value function. Let ϵ be a uniform majoration of that error, i.e. for all k , $\|\epsilon_k\|_\infty \leq \epsilon$. Then

$$\limsup_{k \rightarrow \infty} \|Q^* - Q^{\pi_k}\|_\infty \leq \frac{2\gamma}{(1-\gamma)^2} \epsilon.$$

Proof

Notations and main idea of the proof We will use the following notations:

- $b_k = Q_k - B_{\pi_{k+1}} Q_k$ is the Bellman error,
- $d_k = Q^* - (Q_k - \epsilon_k)$ is the difference between the optimal value function and the Q_k iterate (before error),
- $s_k = Q_k - \epsilon_k - Q^{\pi_k}$ is the difference between the Q_k iterate (before error) and the (true) value of the policy π_k ,
- $\beta = \sum_{n \geq 1} \lambda_n \gamma^n$ (note that $0 \leq \beta \leq \gamma$).

The distance between the value of the optimal policy and the value of the current policy can be formulated as

$$\begin{aligned}
\|Q^* - Q^{\pi_k}\|_\infty &= \max(Q^* - Q^{\pi_k}) \\
&= \max(Q^* - Q_k + \epsilon_k + Q_k - \epsilon_k - Q^{\pi_k}) \\
&= \max(d_k + s_k) \\
&\leq \max d_k + \max s_k
\end{aligned} \tag{1}$$

The idea of the proof is to compute upper bounds on d_k and s_k . As we will see, the bounds we will obtain will both depend on an upper on the Bellman error b_k , that we derive first.

An upper bound on the Bellman error b_k : As π_{k+1} is the greedy policy with respect to Q_k , we have $B_{\pi_k} Q_k \leq B_{\pi_{k+1}} Q_k$, which allows us to write

$$\begin{aligned}
b_k &= Q_k - B_{\pi_{k+1}} Q_k \\
&= Q_k - B_{\pi_k} Q_k + B_{\pi_k} Q_k - B_{\pi_{k+1}} Q_k \\
&\leq Q_k - B_{\pi_k} Q_k \\
&= (Q_k - \epsilon_k + \epsilon_k) - B_{\pi_k} (Q_k - \epsilon_k + \epsilon_k) \\
&= (Q_k - \epsilon_k) - B_{\pi_k} (Q_k - \epsilon_k) + \epsilon_k - \gamma P_{\pi_k} \epsilon_k \\
&= \sum_{n \geq 1} \lambda_n [(B_{\pi_k})^n Q_{k-1}] - \sum_{n \geq 1} \lambda_n [(B_{\pi_k})^{n+1} Q_{k-1}] + (I - \gamma P_{\pi_k}) \epsilon_k \\
&= \sum_{n \geq 1} \lambda_n [(B_{\pi_k})^n Q_{k-1}] - (B_{\pi_k})^{n+1} Q_{k-1} + (I - \gamma P_{\pi_k}) \epsilon_k \\
&= \sum_{n \geq 1} \lambda_n (\gamma P_{\pi_k})^n (Q_{k-1} - B_{\pi_k} Q_{k-1}) + (I - \gamma P_{\pi_k}) \epsilon_k \\
&= \sum_{n \geq 1} \lambda_n (\gamma P_{\pi_k})^n b_{k-1} + (I - \gamma P_{\pi_k}) \epsilon_k.
\end{aligned}$$

By using the fact that P_{π_k} is a stochastic matrix, we have

$$\max b_k \leq \sum_{n \geq 1} \lambda_n \gamma^n \max b_{k-1} + (1 + \gamma) \epsilon = \beta \max b_{k-1} + (1 + \gamma) \epsilon.$$

We then deduce by induction that

$$\max b_k \leq \sum_{j=0}^{k-1} \beta^j (1 + \gamma) \epsilon + \beta^k \max b_0 = \frac{1 + \gamma}{1 - \beta} \epsilon + O(\gamma^k). \tag{2}$$

An upper bound on d_k : Let us now consider the d_k term and its evolution.

$$\begin{aligned}
d_{k+1} &= Q^* - (Q_{k+1} - \epsilon_{k+1}) \\
&= Q^* - \sum_{n \geq 1} \lambda_n (B_{\pi_{k+1}})^n Q_k \\
&= \sum_{n \geq 1} \lambda_n [Q^* - (B_{\pi_{k+1}})^n Q_k].
\end{aligned} \tag{3}$$

Since π_{k+1} is the greedy policy with respect to Q_k , we have $B_{\pi^*} Q_k \leq B_{\pi_{k+1}} Q_k$. Therefore

$$\begin{aligned}
& Q^* - (B_{\pi_{k+1}})^n Q_k \\
&= B_{\pi^*} Q^* - B_{\pi^*} Q_k + B_{\pi^*} Q_k - B_{\pi_{k+1}} Q_k + B_{\pi_{k+1}} Q_k - \\
&\quad - (B_{\pi_{k+1}})^2 Q_k + (B_{\pi_{k+1}})^2 Q_k - \dots + (B_{\pi_{k+1}})^{n-1} Q_k - (B_{\pi_{k+1}})^n Q_k \\
&\leq B_{\pi^*} Q^* - B_{\pi^*} Q_k + \gamma P_{\pi_{k+1}} (Q_k - B_{\pi_{k+1}} Q_k) + \\
&\quad + (\gamma P_{\pi_{k+1}})^2 (Q_k - B_{\pi_{k+1}} Q_k) + \dots + (\gamma P_{\pi_{k+1}})^{n-1} (Q_k - B_{\pi_{k+1}} Q_k) \\
&= \gamma P_{\pi^*} (Q^* - Q_k) + \\
&\quad + [\gamma P_{\pi_{k+1}} + (\gamma P_{\pi_{k+1}})^2 + \dots + (\gamma P_{\pi_{k+1}})^{n-1}] (Q_k - B_{\pi_{k+1}} Q_k) \\
&= \gamma P_{\pi^*} (Q^* - (Q_k - \epsilon_k)) - \gamma P_{\pi^*} \epsilon_k + \\
&\quad + [\gamma P_{\pi_{k+1}} + (\gamma P_{\pi_{k+1}})^2 + \dots + (\gamma P_{\pi_{k+1}})^{n-1}] (Q_k - B_{\pi_{k+1}} Q_k) \\
&= \gamma P_{\pi^*} d_k - \gamma P_{\pi^*} \epsilon_k + [\gamma P_{\pi_{k+1}} + (\gamma P_{\pi_{k+1}})^2 + \dots + (\gamma P_{\pi_{k+1}})^{n-1}] b_k.
\end{aligned}$$

As P_{π^*} and $P_{\pi_{k+1}}$ are stochastic matrices, we deduce

$$\begin{aligned}
\max[Q^* - (B_{\pi_{k+1}})^n Q_k] &\leq \gamma \max d_k + \gamma \epsilon + (\gamma + \gamma^2 + \dots + \gamma^{n-1}) \max b_k \\
&= \gamma \max d_k + \gamma \epsilon + \frac{\gamma - \gamma^n}{1 - \gamma} \max b_k.
\end{aligned}$$

By using Equation 3, we obtain the following induction on $\max d_k$:

$$\max d_{k+1} \leq \gamma \max d_k + \gamma \epsilon + \sum_{n \geq 1} \lambda_n \left[\frac{\gamma - \gamma^n}{1 - \gamma} \max b_k \right].$$

With the help of the Bellman error upper bound obtained earlier (Equation 2) we obtain

$$\begin{aligned}
\max d_{k+1} &\leq \gamma \max d_k + \gamma \epsilon + \sum_{n \geq 1} \lambda_n \left[\frac{\gamma - \gamma^n}{(1 - \gamma)(1 - \beta)} \right] (1 + \gamma) \epsilon + O(\gamma^k) \\
&= \gamma \max d_k + \gamma \epsilon + \frac{\gamma - \beta}{(1 - \gamma)(1 - \beta)} (1 + \gamma) \epsilon + O(\gamma^k)
\end{aligned}$$

which gives, by taking the limit superior,

$$\limsup_{k \rightarrow \infty} \max d_k \leq \frac{\gamma}{1 - \gamma} \epsilon + \left[\frac{\gamma - \beta}{(1 - \gamma)^2 (1 - \beta)} \right] (1 + \gamma) \epsilon. \quad (4)$$

An upper bound on s_k : Let us now consider the s_k term from Equation 1:

$$\begin{aligned}
s_{k+1} &= Q_{k+1} - \epsilon_{k+1} - Q^{\pi_{k+1}} \\
&= \sum_{n \geq 1} \lambda_n [(B_{\pi_{k+1}})^n Q_k] - (B_{\pi_{k+1}})^\infty Q_k \\
&= \sum_{n \geq 1} \lambda_n [(B_{\pi_{k+1}})^n Q_k - (B_{\pi_{k+1}})^\infty Q_k]. \quad (5)
\end{aligned}$$

It can be seen that

$$\begin{aligned}
& (B_{\pi_{k+1}})^n Q_k - (B_{\pi_{k+1}})^\infty Q_k \\
&= (B_{\pi_{k+1}})^n Q_k - (B_{\pi_{k+1}})^{n+1} Q_k + (B_{\pi_{k+1}})^{n+1} Q_k - (B_{\pi_{k+1}})^{n+2} Q_k + \dots \\
&= (\gamma P_{\pi_{k+1}})^n (Q_k - B_{\pi_{k+1}} Q_k) + (\gamma P_{\pi_{k+1}})^{n+1} (Q_k - B_{\pi_{k+1}} Q_k) + \dots \\
&= (\gamma P_{\pi_{k+1}})^n [I + \gamma P_{\pi_{k+1}} + (\gamma P_{\pi_{k+1}})^2 + \dots] b_k.
\end{aligned}$$

As above, by using the stochasticity of $P_{\pi_{k+1}}$, we obtain

$$\max[(B_{\pi_{k+1}})^n Q_k - (B_{\pi_{k+1}})^\infty Q_k] \leq \gamma^n (1 + \gamma + \gamma^2 + \dots) \max b_k = \frac{\gamma^n}{1 - \gamma} \max b_k.$$

By using Equation 5, we obtain an upper bound on $\max s_{k+1}$:

$$\max s_{k+1} \leq \frac{1}{1 - \gamma} \left[\sum_{n \geq 1} \lambda_n \gamma^n \max b_k \right].$$

With the help of the Bellman error upper bound (Equation 2) and by taking the limit superior, we have

$$\limsup_{k \rightarrow \infty} \max s_k \leq \frac{1}{1 - \gamma} \left(\sum_{m \geq 1} \lambda_m \gamma^m \frac{1 + \gamma}{1 - \beta} \epsilon \right) = \frac{\beta}{(1 - \gamma)(1 - \beta)} (1 + \gamma) \epsilon. \quad (6)$$

Conclusion of the proof Finally, let us get back to Equation 1 and use the upper bounds we just derived for d_k (Equation 4) and s_k (Equation 6):

$$\begin{aligned}
\limsup_{k \rightarrow \infty} \|Q^* - Q^{\pi_k}\|_\infty &\leq \limsup_{k \rightarrow \infty} \max d_k + \limsup_{k \rightarrow \infty} \max s_k \\
&= \frac{\gamma}{1 - \gamma} \epsilon + \left[\frac{\gamma - \beta}{(1 - \gamma)^2 (1 - \beta)} + \frac{\beta}{(1 - \gamma)(1 - \beta)} \right] (1 + \gamma) \epsilon. \\
&= \frac{\gamma}{1 - \gamma} \epsilon + \left[\frac{\gamma - \beta + (1 - \gamma)\beta}{(1 - \gamma)^2 (1 - \beta)} \right] (1 + \gamma) \epsilon. \\
&= \frac{\gamma}{1 - \gamma} \epsilon + \left[\frac{\gamma}{(1 - \gamma)^2} \right] (1 + \gamma) \epsilon. \\
&= \frac{\gamma(1 - \gamma) + \gamma(1 + \gamma)}{(1 - \gamma)^2} \epsilon \\
&= \frac{2\gamma}{(1 - \gamma)^2} \epsilon. \quad \blacksquare
\end{aligned}$$

References

Thiery, C. and B. Scherrer (2010). Least-squares λ policy iteration: Bias-variance trade-off in control problems. In *ICML'10: Proceedings of the 27th Annual International Conference on Machine Learning*.